Psychopathology, psychosocial factors and obesity Psicopatologia, fattori psicosociali e obesità

FEDERICA PINNA¹, CLAUDIA SARDU², WALTER ORRÙ^{1,3}, FERNANDA VELLUZZI⁴, ANDREA LOVISELLI⁴, PAOLO CONTU², BERNARDO CARPINIELLO¹ E-mail: bcarpini@iol.it

¹Department of Public Health, Clinical and Molecular Medicine, Unit of Psychiatry, University of Cagliari, Italy ²Department of Public Health, Clinical and Molecular Medicine, Unit of Epidemiology, University of Cagliari, Italy ³H. Shultz Institute, Cagliari, Italy ⁴Centre for Diagnosis and Treatment of Obesity, University Hospital, Cagliari, Italy

SUMMARY. Aims. The aims of this study were to evaluate the association between obesity and socio-demographic and psychopathological variables in a clinical sample of patients referred to a center for the diagnosis and treatment of obesity, compared with a homogeneous sample of normal-weight subjects. **Methods.** In the context of a research project regarding obesity and psychopathology, a multivariate logistic regression analysis was used to examine the association between obesity and the demographic and clinical variables, on the basis of the data set of a consecutive sample of 293 obese patients (48 males, 245 females, mean age: 45.41 ± 13.55 , mean body mass index [BMI] 35.6 ± 6.2) compared with a control group of 293 non-obese subjects (48 males, 245 females, mean age: 45.66 ± 13.86 , mean BMI 21.8 ±2.06). All subjects were assessed by structured clinical interview, the Structured Clinical Interview for Diagnosis for axis I DSM-IV (SCID-I) and for axis II DSM-IV (SCID-II). **Results.** Multivariate statistical analysis showed that the status of housewife and the presence of lifetime axis I and II psychiatric diagnosis in general, and of depressive, anxiety, eating and some personality disorders in particular, significantly increased the likelihood of being overweight/obese. The likelihood for different combinations of risk factors considered. **Conclusions.** The presence of an axis I and/or II diagnosis and risk factor, to a value of 86.7% for those exposed to all risk factors considered. **Conclusions.** The presence of an axis I and/or II diagnosis increases this likelihood. Even taking into account the limits of the study, in particular of its cross-sectional nature, these findings may have important implications in both prevention and treatment of obesity.

KEY WORDS: obesity, overweight, mental disorders, depression, anxiety, eating disorders, personality disorders, comorbidity, psychosocial factors.

RIASSUNTO. Scopi. Gli obiettivi di questo studio sono stati la valutazione dell'associazione tra obesità, variabili socio-demografiche e psicopatologiche in un campione clinico di pazienti afferenti a un centro per la diagnosi e il trattamento dell'obesità, confrontati con un campione omogeneo di controllo costituito da soggetti di peso normale. Metodi. Nel contesto di un progetto di ricerca riguardante obesità e psicopatologia, è stata applicata una regressione logistica multivariata per esaminare l'associazione tra obesità e variabili demografiche e cliniche relative a un campione consecutivo di 293 pazienti obesi (48 maschi, 245 femmine, età media: 45,41±13,55, indice di massa corporea [BMI] 35.6±6.2), confrontato con un gruppo di controllo omogeneo appaiato di 293 soggetti non obesi (48 maschi, 245 femmine, età media 45,66±13,86, BMI medio 21,8±2,06). Tutti i soggetti sono stati valutati mediante un'intervista clinica strutturata, la Structured Clinical Interview per la diagnosi di asse I del DSM-IV (SCID-I) e per l'asse II DSM-IV (SCID-II). Risultati. L'analisi statistica multivariata ha dimostrato che lo status di casalinga e la presenza "lifetime" di disturbi di asse I e II e in generale della presenza di una qualsiasi diagnosi psichiatrica, e, in particolare, delle diagnosi di disturbo depressivo, d'ansia, del comportamento alimentare e di alcuni disturbi di personalità, sono associate in modo significativo a una maggiore probabilità di essere in sovrappeso oppure obesi. Le probabilità di essere sovrappeso/obesi si modificano in rapporto alle diverse combinazioni di fattori di rischio, aumentando da un valore di 32,3% per un individuo non esposto ad alcun fattore di rischio, a un valore di 86,7% per i soggetti esposti a tutti i fattori di rischio considerati. Conclusioni. La presenza di una diagnosi di asse I e/o II e lo stato di casalinga è indipendentemente associata a un maggiore rischio di sovrappeso/obesità. L'interazione di questi fattori aumenta questo rischio. Pur tenendo conto dei limiti dovuti alla natura trasversale dello studio, questi risultati sono collegati a implicazioni importanti sia ai fini della prevenzione sia del trattamento dell'obesità.

PAROLE CHIAVE: obesità, sovrappeso, disturbi mentali, depressione, ansia, disturbi alimentari, disturbi di personalità, comorbilità, fattori psicosociali.

INTRODUCTION

Overweight and obesity are a public health problem featuring a constant upwards trend^{1,2} and represent a relevant risk factor for a number of other medical conditions associated with increased rates of morbidity and mortality^{1.4}. The multifactorial etiology of obesity involves complex interactions between genetic, hormonal and social-environmental

Psychopathology, psychosocial factors and obesity

factors, including sedentary lifestyle and poor eating habits^{1,2}.

Several other factors, including age, gender, socio-cultural, economic and marital status and job categories have been associated with the development of overweight/obesity⁵⁻⁷. Psychosocial factors occurring in childhood, such as episodes of violence and abuse, poor parental support and academic difficulties, appear to produce a significant impact in determining the development of overweight/obesity in adolescence and adulthood^{8,9}. Moreover, a relevant number of cross-sectional studies, the majority of which focused on depression, have revealed a significant association of psychopathology with overweight/obesity. Indeed, a meta-analysis of 17 crosssectional population studies has shown that depression is associated with an increased risk of becoming obese, an association found in women but not in men¹⁰. Moreover, an association between obesity and anxiety disorders emerged from a recent review and meta-analysis of population-based studies, largely based upon a cross-sectional design¹¹. Most of the studies show a positive relationship with panic disorder, mainly in women, and with specific phobia and social phobia, although the role of subtypes of anxiety disorders and obesity severity remains to be fully clarified¹². Similarly, a strong association is reported in literature between eating disorders (ED), particularly binge eating disorder (BED) and obesity¹³. Finally, data currently available derived from cross-sectional studies show a high overall prevalence of personality disorders in obese subjects¹⁴⁻¹⁷. As cross-sectional studies do not allow any inference to be made as to the direction of causality (i.e. mental disorder may be the cause or consequence of obesity), more information is provided by longitudinal studies, again prevalently focused on depression. Indeed, the presence of depressive symptoms during childhood and adolescence has been revealed as one of the most frequently identified risk factors in relation to the development of obesity in adulthood, particularly among women^{8,10,18}. However, a review and meta-analysis of longitudinal studies has demonstrated a reciprocal relationship between obesity and depression, with important implications in terms of prevention, early diagnosis and treatment¹⁹. Indeed, a condition of overweight/obesity was found to increase the risk of depression, whilst on the other hand depression was found to be predictive of later development of obesity, with a 58% increased risk of becoming obese¹⁹. With regard to anxiety, a recent prospective cohort study showed that symptoms of anxiety, on a par with symptoms of depression, were associated with a greater change in weight and increased incidence of obesity in both men and women²⁰. Several longitudinal studies have demonstrated binge eating as a risk factor for overweight/obesity²¹⁻²⁶, although a study of adolescent girls failed to identify this association²⁷. Pertaining to personality psychopathology, a longitudinal study showed that aggressive personality traits and sociopathy were positively associated with the development of overweight²³, with another study revealing a positive relationship between some specific personality disorders in early adulthood and an increased risk of problems related to eating habits and weight in later life²⁸. Overall, a large body of literature, including large com-munity studies^{10,11,17,29,30} has reported a significant association between specific psychopathological conditions, mainly depression, and overweight/obesity, with a bi-directionality of causal relationship. Starting from these premises and considering the relative paucity of studies of clinical populations of obese patients conducted to verify diagnosis by means of structured psychiatric interviews, a study program on obesity and psychopathology was set up several years ago, the results of which have been previously reported elsewhere³¹⁻³⁵. In a first cross-sectional study of 150 consecutive obese patients attending a university center for the diagnosis and treatment of obesity we demonstrated a very high lifetime prevalence of axis I (58%) and axis II (28%) disorders according to DSM-IV-TR³¹. Prevalence rates were significantly higher than those found in the corresponding general population across all principal classes of common mental disorders (mood and anxiety disorders) and among ED, with depressive disorders (DD) (26%) and BED (16%) being the single most widely represented diagnoses. In a subsequent casecontrolled study³², the principal aim of which was to investigate quality of life in relation to psychopathology, a larger sample of obese patients (n=293) was studied and compared to a sample of normal controls (n=293) matched according to gender, age, marital status and education; in this study a significantly higher odds of both axis I and II disorders was detected among obese subjects, in particular anxiety and mood disorders, with particular reference to major depression (MD), and BED³². In the present paper we report the results of a further analysis of the data set obtained in the latter study, for the purpose of examining, by means of multivariate analysis, the association between obesity and psychiatric disorders, taking into account social variables, potential confounding effects and the interaction of factors.

MATERIALS AND METHODS

Sample

An unselected sample of consecutive patients referred to a University Center for the diagnosis and treatment of obesity that treats approx. 450 subjects a year was assessed. The center provides a comprehensive medical and psychological evaluation with a multidisciplinary approach, including drug treatment, dietary and psycho-educational interventions tailored to individual needs. When necessary, psychological support or psychiatric intervention was made available. Of the 298 subjects contacted over a 9-month period, only five refused to take part in the study. The final study group was made up of 293 subjects. As a control group of normalweight subjects, we used a convenience sample of university students and their families made up of 293 individuals, matched as much as possible for sex, age, marital status and education. Sample and controls were prevalently constituted by females (48 males, 245 females, both in cases and controls), married (respectively 199, 67.9% and 198, 67.6%), with a high level of education (high school or graduates were respectively 155, 52.9% and 156, 53.2%), without any difference from a statistical point of view; as far as occupation was concerned, a higher number of employed was found among controls (167,57% vs 127,43.4% of cases) and of housewives among cases (92, 31.4% vs 64, 21.8% of controls) (p<.001). As expected, mean BMI (cases=35.60+/-6.20, controls=21.83+/-2.06) and average waist circumference (cases=112.17+/-13.94 cm, controls=78.38+/-9.58 cm) were both significantly higher among cases than controls (respectively, t=36.047, df=584, p<0.001, t=34.173, df=584, p<0.001). Among cases approx. 16.8% (n=50) of subjects were overweight; obese subjects were distributed in classes as follows: class I 33.2% (n=97); class II 29.1% (n=85), class III 20.9% (n=292).

Pinna F et al.

Methods

Socio-demographic data, medical history and clinical data were collected by means of a structured interview purpose-developed for the study.

After signing an informed consent form, all patients and controls underwent a comprehensive psychiatric evaluation using the Structured Clinical Interview for Diagnosis for axis I DSM-IV (SCID-I Research Version)³⁶, and the interview for the Structured Clinical Diagnosis of axis II of DSM-IV (SCID-II)³⁷.

Interviews were conducted by residents in psychiatry trained in the use of the instrument by a trained specialist (FP), the interrater reliability, assessed using Cohen's K before starting the study, was not less than 0.80.

Anthropometric measurements were taken using standard calibrated instruments. Height (m) was measured using a wallmounted stadiometer; weight (kg) using electronic scales with an upper weight limit of 300 kg. Body Mass Index (BMI) was calculated as Weight/Height².

Statistical analysis

Differences between continuous variables were calculated by means Student's t test for unpaired data; comparison of categorical variables was performed by means of chi square test. Basing upon data of preliminary univariate analyses previously published in another paper of our group³², in order to assess the combined effects of different variables on the risk of obesity, a multivariate logistic regression was performed, considering obesity vs normal weight status as dependent variables, while sociodemographic variables and the presence of an axis I or axis II diagnosis were considered as independent variables. The dependent variable was analyzed as a function of each independent variable and their second order interactions. The backward-stepwise method was adopted, in which all variables and interactions were initially inserted in the model based on the available clinical and epidemiological knowledge; subsequently, non-significant interactions (p>0.20) and variables (p>0.05) were consecutively eliminated by means of a step by step procedure starting from the saturated model applied to perform logistic regression. The association of each independent variable with the dependent variable was expressed by means of OR and its 95% confidence interval. The probability of being obese as a function of the different combinations of risk factors identified was calculated using the parameters obtained with the logistic model. A further logistic regression was used to identify the specific diagnosis of axis I and axis II associated with obesity. This second analysis considered the status of being overweight vs. normal weight as dependent variable and a number of diagnoses (bipolar disorder, major depressive disorder or other unipolar disorders, anxiety disorders, eating disorders, substance use disorders, psychotic disorders, avoidant personality disorder, obsessive compulsive personality disorder, passive aggressive personality disorder, depressive personality disorder, paranoid personality disorder, schizoid personality disorder, histrionic personality disorder, narcissistic personality disorder, borderline personality disorder, antisocial personality disorder, personality disorder not otherwise specified [NOS]), as independent variables, in addition to professional status.

RESULTS

Univariate analyses demonstrated that obesity was associated with a significant lifetime major risk both for axis I (OR=3.47, p=0.000) and axis II disorders $(OR=2.27, p=0.000)^{32}$.

No statistically significant difference was found in distribution of subjects according to overweight/obesity classes and psychiatric diagnosis (any axis I, any axis II disorders or any specific disorder); with only two exceptions: MD and obsessive compulsive disorder (OCD). Indeed, patients affected by MD (n=59) included a significantly higher proportion of overweight (20.3%), class I (35.6%) and II (37.3%) obese subjects and a lower proportion of class III (6.8%) subjects compared to patients affected by any other disorder (n=234), who were diagnosed as follows: overweight=15.9%, class I=32.6%, class II=27%, class III=24.5% (chi square test=9.430, gl=3, p=0.024). Patients affected by OCD (n=4) were characterized by a significantly different distribution, with all 4 subjects classified as obesity class I (100%), compared to patients affected by any other disorder (n=288) who were classified as follows: overweight=17%, class I=32.3%, class II=29.5%, class III=21.2% (chi square test=8.153, gl=3, p=0.043).

Multivariate logistic regression showed a significant effect of professional status (housewife) and both axis I and axis II diagnosis (Table 1). The final model indicated that individuals with a diagnosis of axis I run a 2.7 times greater likelihood of obesity compared with subjects without axis I disorders (95% CI 1.20 to 2.63). Individuals with an axis II diagnosis are at a 2.80 times greater likelihood of obesity compared with subjects with no axis II diagnosis. Housewives showed an approx. 1.8 times greater likelihood of obesity than other professional statuses considered. The model showed no significant interactions, so that the likelihood of obesity in presence of more than one risk factor is the product of the individual odds ratios. The likelihood of being obese for different combinations of risk factors increased from a value of 32.3% for an individual not exposed to any risk factor, to a value of 86.7% for those exposed to all risk factors considered (Table 2).

Table 1. Multivariate Logistic Regression analysis: significant associations between obesity (dependent variable) and clinical/psychosocial variables (independent variables)

	р	OR	CI 95%
Housewife	0.004	1.77	1.20-2.63
Lifetime axis I diagnosis	0.0000001	2.75	1.89-3.98
Axis II diagnosis	0.0001	2.80	1.70-4.62

Table 2. Likeliho	od of	being	obese	for	different	combinations	of
risk factors		-					

Risk factors	Likelihood of being obese
No risk factor	32.3%
Housewife	45.8%
Lifetime axis I diagnosis	56.7%
Axis II diagnosis	57.2%
Housewife+axis I diagnosis	69.9%
Housewife+axis II diagnosis	70.3%
Axis I+axis II diagnosis	78.6%
Housewife+axis I+axis II diagnosis	86.7%

Psychopathology, psychosocial factors and obesity

When logistic regression was used to identify the specific axis I and axis II diagnoses associated with obesity, a higher likelihood associated with the status of housewife was again confirmed, together with being affected by major depression (OR=2.17), anxiety disorder (OR=2.35), eating disorder (OR=12.92), avoidant disorder (OR=10.07), paranoid disorder (OR=11.43), personality disorder NOS (OR=1.96) (Table 3).

DISCUSSION

Before discussion data from this study, it should be underlined that findings should be considered in the light of several major limitations. First, the sample investigated comprised individuals seeking weight loss treatment and should therefore not be considered representative of obese subjects living in the community or of obese people contacting other clinical settings for reasons other than their overweight; second, the sample examined was largely made up of females, a finding consistent with other studies investigating clinical samples^{40,41}, likely reflecting a more pronounced propensity of women to seek assistance for weight related problems; third, mental disorders were evaluated only by means of structured clinical interviews, without using self-reported instruments; fourth, the cross-sectional nature of this study prevents the drawing of any conclusion as to the direction of causality in the association between mental disorders and obesity.

Data available in the literature show consistently higher prevalence rates of mental disorders in clinical³⁸⁻⁴³ samples of obese patients. The association of mental health and obesity may be related to an interplay of a wide range of individual (such as gender and age) and contextual factors (such race, ethnicity and culture)44,45. Moreover, several biological factors have been proposed to explain the relationship between mental disorders and obesity with significant implications for the treatment⁴⁶.

Preliminary results of this study project, based upon univariate analyses³², showed a significantly increased odds of any lifetime axis I and axis II disorders, axis II disorders or their comorbidity to be associated with obesity; in particular, among axis I disorders, mood disorders (including MD, BD), anxiety disorders, ED were strongly associated with over-

Table 3. Multivariate Logistic Regression analysis: significant associations between obesity (dependent variable) and categories of mental disorders (independent variables)

p OR CI 95% Housewife 0.003 1.83 1.23-2.73 Major depressive disorder or other unipolar disorders 0.001 2.17 1.35-3.50 Anxiety disorders 0.0003 2.35 1.48-3.73 Eating disorders 0.0003 12.92 3.88-42.97 Avoidant personality disorder 0.03 10.07 1.18-85.72 Paranoid personality disorder 0.02 11.43 1.37-95.37 Personality disorder NOS 0.03 1.96 1.06-3.62				
Housewife0.0031.831.23-2.73Major depressive disorder or other unipolar disorders0.0012.171.35-3.50Anxiety disorders0.00032.351.48-3.73Eating disorders0.000312.923.88-42.97Avoidant personality disorder0.0310.071.18-85.72Paranoid personality disorder0.0211.431.37-95.37Personality disorder NOS0.031.961.06-3.62		р	OR	CI 95%
Major depressive disorder or other unipolar disorders0.0012.171.35-3.50Anxiety disorders0.00032.351.48-3.73Eating disorders0.0000312.923.88-42.97Avoidant personality disorder0.0310.071.18-85.72Paranoid personality disorder0.0211.431.37-95.37Personality disorder NOS0.031.961.06-3.62	Housewife	0.003	1.83	1.23-2.73
Anxiety disorders 0.0003 2.35 1.48-3.73 Eating disorders 0.00003 12.92 3.88-42.97 Avoidant personality disorder 0.03 10.07 1.18-85.72 Paranoid personality disorder 0.02 11.43 1.37-95.37 Personality disorder NOS 0.03 1.96 1.06-3.62	Major depressive disorder or other unipolar disorders	0.001	2.17	1.35-3.50
Eating disorders 0.00003 12.92 3.88-42.97 Avoidant personality disorder 0.03 10.07 1.18-85.72 Paranoid personality disorder 0.02 11.43 1.37-95.37 Personality disorder NOS 0.03 1.96 1.06-3.62	Anxiety disorders	0.0003	2.35	1.48-3.73
Avoidant personality disorder0.0310.071.18-85.72Paranoid personality disorder0.0211.431.37-95.37Personality disorder NOS0.031.961.06-3.62	Eating disorders	0.00003	12.92	3.88-42.97
Paranoid personality disorder0.0211.431.37-95.37Personality disorder NOS0.031.961.06-3.62	Avoidant personality disorder	0.03	10.07	1.18-85.72
Personality disorder NOS0.031.961.06-3.62	Paranoid personality disorder	0.02	11.43	1.37-95.37
	Personality disorder NOS	0.03	1.96	1.06-3.62

weight. Basing on these previous findings, in this paper we re-examined the association between obesity and psychiatric disorders by means of multivariate analyses, taking into account psychosocial variables and the potential confounding effects and interaction of factors.

In particular, multivariate analysis revealed how being an housewife was the only socio-demographic condition associated to an increased likelihood of overweight/obesity. This finding is congruent with the limited data present in literature showing a higher prevalence of obesity among housewives compared to other workers^{7,47}. Numerous factors have been hypothesized as underlying these findings, including low socio-economic and/or cultural status, unhealthy dietary habits and insufficient physical activity, given that family and domestic duties may represent a limitation with respect to a commitment to regular physical activity⁷. However, the fact that physical activity alone has not proved to be sufficient to obtain and maintain a substantial weight loss should be considered47.

Multivariate analysis confirmed that being affected by a lifetime axis I diagnosis, in particular MD disorder or other unipolar mood disorders, anxiety disorders and eating disorders (BED) is associated with an increased likelihood of overweight/obesity. Likewise, being affected by an axis II personality disorder, in particular avoidant, paranoid, and not otherwise specified disorder, is also correlated with an increased odds of overweight/obesity. These results are largely in accordance with data currently available in the literature with regard to the association of psychopathology and overweight/obesity. Accordingly, the findings reported in longitudinal studies have demonstrated that depressive symptoms during childhood and adolescence are associated with an increased risk of development of obesity in adulthood^{8,19,48}. Several putative reasons have been put forward to explain this increased risk of overweight/obesity among depressed subjects. Unhealthy lifestyles, both in terms of insufficient physical activity and eating habits have been implicated, particularly in atypical depression, frequently associated with binge eating and overeating^{19,49-51}. Antidepressant treatments may lead to a weight gain in subjects affected by depression or other mental disorders¹⁹. Dysfunction of biological systems implicated in both depression and obesity, such as the hypothalamic-pituitary-adrenal (HPA) axis, the central monoaminergic systems, leptin, the immune systems and glucose and lipid metabolism may interact, thus contributing towards explaining such a frequent association between depression and overweight/obesity⁵²⁻⁵⁶. In agreement with our results, literature data emphasizing the association between obesity and anxiety disorders reveal an overall positive relationship^{11,12}. Data from a prospective study showed an association between anxiety disorders in childhood and adolescence and an excess of weight in subsequent years, although at a statistically significant level only for females⁵⁷. A recent longitudinal population study demonstrated that both male and female subjects who received a diagnosis of anxiety disorder during infancy featured a higher BMI than controls in the following years⁴⁸. The risk of weight gain related to an anxiety disorder has itself been attributed to different factors such as a possible dysregulation of the HPA axis, dysfunctional eating habits due to symptoms such as increased appetite and/or craving for sweet or fat foods, the presence a comorbid chronic condition (e.g., asthma) with a significant

Pinna F et al.

impact on functioning, resulting in decline in physical activity and excess weight¹¹. In addition, anxiety disorders may indirectly be related to an increased odds of overweight/obesity though genetic and hormonal factors, traumatic events in childhood, personality dysfunctional traits, comorbidity with other psychiatric and somatic conditions¹¹. In our study binge eating disorder was seen to be strongly associated with an increased likelihood of overweight/obesity, a result in agreement with the findings obtained in the majority of longitudinal studies, showing a positive association between binge eating and subsequent weight gain²¹⁻²⁶. However, it should be underlined that even a study reporting a negative association has been published²⁷. Uncontrolled overeating is viewed as one of the main explanations for the weight gain frequently observed among patients affected by BED; moreover, a common genetic factor shared by patients affected by obesity and binge eating disorder was hypothesized^{58,59}. Our data lend further support to the hypothesis that some personality disorders may contribute towards the development of weight problems, as shown by previous cross-sectional studies showing a higher BMI in subjects with a cluster A or B personality disorders^{60,61}, higher percentages of cluster A, B, and C personality disorders in subjects with severe obesity⁶⁰, an increased odds of avoidant and antisocial personality disorder in obese women with extreme obesity¹⁷, and a positive association between overweight/obesity and paranoid, antisocial and avoidant personality disorders among female patients¹⁵. Moreover, a longitudinal community study has demonstrated an association between several personality disorders, namely schizoid, schizotypal and antisocial disorders in early adulthood, and a major risk of weight problems or obesity in later life²⁸. The reason for this association is unknown, although some behavioral factors may partly explain these findings, such as a tendency towards marked social avoidance and the consequent development of a sedentary lifestyle in individuals affected by avoidant and paranoid personalities, while impulsiveness and the related tendency to loss of control over eating habits could be taken into account for individuals with antisocial and borderline personality disorders¹⁵.

Some authors have claimed that the frequent association between cluster A personality disorders and obesity may be due to a common dopaminergic dysfunction, although this hypothesis fails to explain the gender differences observed in the association between obesity and these personality disorders reported in some studies^{15,17}. One of the main findings of the present study is the apparent interactive role of social and psychopathological factors in increasing the odds of obesity. Indeed, the probability of being overweight/obese rises approx. to 46% for a housewife and to approx 57% for a person affected by an axis I or II disorder; this probability increases to approx. 70% for a housewife affected by an axis I or II disorder, becoming extremely high for subjects affected by an axis I and II disorder (approx. 79%) and for housewives featuring comorbidity with an axis I and II disorder (approx. 87%). In other words, our study demonstrates how the presence of an axis I and II disorder according to DSM-IV-TR, and being a housewife are independently associated to a higher odds of being obese, highlighting how the interaction of these factors further increases this probability. Therefore, the evidence provided by this study appears to be of interest not only from a theoretical, but also from a practical point of view, particularly in view of the potential use in identifying subjects at risk of developing obesity. Although our data are congruent with those deriving from longitudinal studies, supporting the possible role of several psychiatric conditions in conjunction with psychosocial variables in developing obesity, as previously acknowledged, the cross-sectional nature of our study prevents the drawing of any conclusion in terms of direction of causality in the association between mental disorders and obesity. Indeed, psychopathology may be not only the cause but even a consequence of obesity, at least as regard to depression¹⁹, a hypothesis further supported by studies regarding remission of depression after surgical treatment of bariatric patients⁶². However, irrespective of the nature of the relationship between obesity and mental disorders (whichever is the prevalent disorder), the data obtained demonstrate that more than half of obese patients seeking treatment for their overweight or obesity are affected by psychiatric comorbidity, a finding which justifies the need for careful psychiatric evaluation and treatment in the context of a multidisciplinary approach.

Conflict of interests: the authors declare they have no competing interests.

Acknowledgements: the authors thank Ms Anne Farmer for her contribution to the translation of the paper into English.

REFERENCES

- Chan RS, Woo J. Prevention of overweight and obesity: how effective is the current public health approach. Int J Environ Res Public Health 2010; 7: 765-83.
- Knight JA. Diseases and disorders associated with excess body weight. Ann Clin Lab Sci 2011; 41: 107-21.
- 3. Brown WV, Fujioka K, Wilson PW, Woodworth KA. Obesity: why be concerned? Am J Med 2009; 122 (4 suppl 1): S4-11.
- Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. BMC Public Health 2009; 9: 88.
- Schoenborn CA, Adams PF, Barnes PM. Body weight status of adults: United States, 1997-98. Adv Data 2002; (330): 1-15.
- La Rosa E, Valensi P, Cohen R, et al. Socioeconomic determinism of obesity in the Seine-Saint-Denis area. Presse Med 2003; 32: 55-60.
- Navadeh S, Sajadi L, Mirzazadeh A, Asgari F, Haghazali M. Housewives' obesity determinant factors in iran; national survey - stepwise approach to surveillance. Iran J Public Health 2011; 40: 87-95.
- Vamosi M, Heitmann BL, Kyvik KO. The relations between an adverse psychological and social environment in childhood and adult obesity of the development: a systematic literature review. Obes Rev 2010; 11: 177-84.
- 9. Foucart J, De Buck C, Verbanck P. Factorial study of the psychopathological factors of morbid obesity in the adolescent. Encephale 2012; 38: 310-7.
- de Wit L, Luppino M, van Straten A, Penninx B, Zitman F, Cuijpers P. Depression and obesity: a meta-analysis of communitybased studies. Psychiatry Res 2010; 178: 230-5.
- 11. Gariepy G, Nitka D, Schmitz N. The association between obesity and anxiety disorders in the population: a systematic review and meta-analysis. Int J Obes (Lond) 2010; 34: 407-19.
- 12. Lykouras L, Michopoulos J. Anxiety disorders and obesity. Psychiatrike 2011; 22: 307-13.

Psychopathology, psychosocial factors and obesity

- Fandiño J, Moreira RO, Preissler C, et al. Impact of binge eating disorder in the psychopathological profile of obese women. Compr Psychiatry 2010; 51: 110-4.
- Sutin AR, Ferrucci L, Zonderman AB, Terracciano A. Personality and obesity across the adult life span. J Pers Soc Psychol 2011; 101: 579-92.
- Mather AA, Cox BJ, Enns MW, Sareen J. Associations between body weight and personality disorders in a nationally representative sample. Psychosom Med 2008; 70: 1012-9.
- López-Pantoja JL, Cabranes JA, Sanchez-Quintero S, et al. Personality profiles between obese and control subjects assessed with five standardized personality scales. Actas Esp Psiquiatr 2012; 40: 266-74.
- 17. Pickering RP, Grant BF, Chou SP, Compton WM. Are overweight, obesity, and extreme obesity associated with psychopathology? Results from the national epidemiologic survey on alcohol and related conditions. J Clin Psychiatry 2007; 68: 998-1009.
- Kalarchian MA, Marcus MD. Psychiatric comorbidity of childhood obesity. Int Rev Psychiatry 2012; 24: 241-6.
- Luppino FS, de Wit LM, Bouvy PF, et al. Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. Arch Gen Psychiatry 2010; 67: 220-9.
- Brumpton B, Langhammer A, Romundstad P, Chen Y, Mai XM. The associations of anxiety and depression symptoms with weight change and incident obesity: The HUNT Study. Int J Obes (Lond) 2013; 37: 1268-74.
- Tanofsky-Kraff M, Cohen ML, Yanovski SZ, et al. A prospective study of psychological predictors of body fat gain among children at high risk for adult obesity. Pediatrics 2006; 117: 1203-9.
- 22. Sonneville KR, Horton NJ, Micali N, et al. Longitudinal associations between binge eating and overeating and adverse outcomes among adolescents and young adults: does loss of control matter? JAMA Pediatr 2013; 167: 149-55.
- Hasler G, Pine DS, Gamma A, et al. The associations between psychopathology and being overweight: a 20-year prospective study. Psychol Med 2004; 34: 1047-57.
- 24. Field AE, Austin SB, Taylor CB, et al. Relation between dieting and weight change among preadolescents and adolescents. Pediatrics 2003; 112: 900-6.
- Stice E, Presnell K, Spangler D. Risk factors for binge eating onset in adolescent girls: a 2-year prospective investigation. Health Psychol 2002; 21: 131-8.
- 26. Stice E, Cameron RP, Killen JD, Hayward C, Taylor CB. Naturalistic weight-reduction efforts prospectively predict growth in relative weight and onset of obesity among female adolescents. J Consult Clin Psychol 1999; 67: 967-74.
- Stice E, Presnell K, Shaw H, Rohde P. Psychological and behavioral risk factors for obesity onset in adolescent girls: a prospective study. J Consult Clin Psychol 2005; 73: 195-202.
- Johnson JG, Cohen P, Kasen S, Brook JS. Personality disorder traits evident by early adulthood and risk for eating and weight problems during middle adulthood. Int J Eat Disord 2006; 39: 184-92.
- 29. Carpenter KM, Hasina DS, Allison DB, Faith MS. Relationships Between Obesity and DSM-IV Major Depressive Disorder, suicide ideation, and suicide attempts: results from a general population study. Am J Public Health 2000; 90: 251-7.
- Simon GE, Von Korff M, Saunders K, et al. Association between obesity and psychiatric disorders in the US adult population. Arch Gen Psychiatry 2006; 63: 824-30.
- Carpiniello B, Pinna F, Pillai G, et al. Obesity and psychopathology. A study of psychiatric comorbidity among patients attending a specialist obesity unit. Epidemiol Psichiatr Soc 2009; 18: 119-27.
- Carpiniello B, Pinna F, Pillai G, et al. Psychiatric comorbidity and quality of life in obese patients. Results from a case-control study. Int J Psychiatry Med 2009; 39: 63-78.
- Carpiniello B, Pinna F, Pili R, Velluzzi F, Loviselli A. Mental disorders in obese patients with and without metabolic syndrome. Int J Psychiatry Med 2011; 42: 369-75.

- Carpiniello B, Pinna F, Velluzzi F, Loviselli A. Mental disorders in patients with metabolic syndrome. The key role of central obesity. Eat Weight Disord 2012; 17: e259-e266.
- Pinna F, Lai L, Pirarba S, et al. Obesity, alexithymia and psychopathology: a case-control study. Eat Weight Disord 2011; 16: e164-e170.
- 36. First MB, Spitzer RL, Williams JBW, Gibbon M. Structured Clinical Interview for DSM-IV Axis I Disorders-Research Version (SCID I, Version 2.0). New York: Biometrics Research Department, New York State Psychiatric Institute, 1996.
- First MB, Gibbon M, Spitzer RL, Williams JBW, Benjamin L. Structured Clinical Interview for DSM-IV Axis II-Personality Disorders (SCID II, Version 2.0). New York: Biometrics Research Department, New York State Psychiatric Institute, 1996.
- Ramacciotti CE, Coli E, Passaglia C, Lacorte M, Pea E, Dell'Osso L. Binge eating disorder: prevalence and psychopathological features in a clinical sample of obese people in Italy. Psychiatry Res 2000; 94: 131-8.
- Vila G, Zipper E, Dabbas M, et al. Mental disorders in obese children and adolescents. Psychosom Med 2004; 66: 387-94.
- Lin HY, Huang CK, Tai CM, et al. Psychiatric disorders of patients seeking obesity treatment. BMC Psychiatry 2013; 13: 1.
- Black DW, Goldstein RB, Mason EE. Prevalence of mental disorder in 88 morbidly obese bariatric clinic patients. Am J Psychiatry 1992; 149: 227-34.
- Kalarchian MA, Marcus MD, Levine MD, et al. Psychiatric disorders among bariatric surgery candidates: relationship to obesity and functional health status. Am J Psychiatry 2007; 164: 328-34.
- Rovera C, Curti R, Colombo EM, et al. Psychiatric features before and after intervention: a study of patients affected by severe obesity undergoing adjustable gastric banding. Riv Psichiatr 2013; 48: 393-9.
- Assari S. The link between mental health and obesity: role of individual and contextual factors. Int J Prev Med 2014; 5: 247-9.
- 45. Assari S. Additive effects of anxiety and depression on body mass index among blacks: role of ethnicity and gender. Int Cardiovasc Res J 2014; 8: 44-51.
- 46. Lopresti AL, Drummond PD. Obesity and psychiatric disorders: Commonalities in dysregulated biological pathways and their implications for treatment. Prog Neuropsychopharmacol Biol Psychiatry 2013; 45: 92-9.
- Ersoy C, Imamoglu S. Comparison of the obesity risk and related factors in employed and Unemployed (housewife) urban Premenopausal women. Diabetes Res Clin Pract 2006; 72: 190-6.
- Rofey DL, Kolko RP, Iosif AM, et al. A longitudinal study of childhood depression and anxiety in relation to weight gain. Child Psychiatry Hum Dev 2009; 40: 517-26.
- Preiss K, Brennan L, Clarke D. A systematic review of variables associated with the relationship between obesity and depression. Obes Rev 2013; 14: 906-18.
- Zhao G, Ford ES, Dhingra S, Li C, Strine TW, Mokdad AH. Depression and anxiety among US adults: associations with body mass index. Int J Obes (Lond) 2009; 33: 257-66.
- Breslin FC, Gnam W, Franche RL, Mustard C, Lin E. Depression and activity limitations: examining gender differences in the general population. Soc Psychiatry Psychiatr Epidemiol 2006; 41: 648-55.
- McElroy SL, Kotwal R, Malhotra S, Nelson EB, Keck PE, Nemeroff CB. Are mood disorders and obesity related? A review for the mental health professional. J Clin Psychiatry 2004; 65: 634-51.
- Pennix BW, Milaneschi Y, Lamers F, Vogelzangss N. Understanding the somatic consequences of depression: biological mechanisms and the role of depression symptom profile. BMC Med 2013; 11: 129.
- 54. Björntorp P. Do stress reactions causes abdominal obesity and comorbidities? Obes Rev 2001; 2: 73-86.

Pinna F et al.

- Carroll BJ, Cassidy F, Naftolowitz D, et al. Pathophysiology of hypercortisolism in depression. Acta Psychiatr Scand Suppl 2007; (433): 90-103.
- Abelson JL, Khan S, Liberzon I, Young EA. HPA axis activity in patients with panic disorder: review and synthesis of four studies. Depress Anxiety 2007; 24: 66-76.
- 57. Anderson SE, Cohen P, Naumova EN, Must A. Association of depression and anxiety disorders with weight change in a prospective community-based study of children followed up into adulthood. Arch Pediatr Adolesc Med 2006; 160: 285-91.
- Branson R, Potoczna N, Kral JG, Lentes KU, Hoehe MR, Horber FF. Binge eating as a major phenotype of melanocortin 4 receptor gene mutations. N Engl J Med 2003; 348: 1096-103.
- Hudson JI, Lalonde JK, Berry JM, et al. Binge-eating disorder as a distinct familial phenotype in obese individuals. Arch Gen Psychiatry 2006; 63: 313-9.
- Rosmond R, Eriksson E, Björntorp P. Personality disorders in relation to anthropometric, endocrine and metabolic factors. J Endocrinol Invest 1999; 22: 279-88.
- Rosmond R, Baghei F, Holm G, Björntorp P. Relationships between personality disorders and anthropometry, hormones and metabolism in women. J Endocrinol Invest 2001; 24: 159-65.
- 62. Rovera C, Curti R, Colombo EM, et al. Psychiatric features before and after intervention: a study of patients affected by severe obesity undergoing adjustable gastric banding. Riv Psichiatr 2013; 48: 393-9.