

Studi sperimentali

Paranoia, depression and lack of insight in schizophrenia: a suggestion for a mediation effect

Paranoia, depressione e mancanza di insight nella schizofrenia: suggerimenti per una mediazione

FRANCESCA PACITTI^{1*}, DARIO SERRONE¹, ALESSIA LUCASELLI¹, DALILA TALEVI¹,
ALBERTO COLLAZZONI¹, PAOLO STRATTA², RODOLFO ROSSI³

*E-mail: fpacitti@tin.it

¹Department of Biotechnological and Applied Clinical Sciences, Section of Psychiatry, University of L'Aquila, Italy

²Department of Mental Health, ASL 1, Abruzzo, ITALY

³Department of Systems Medicine, University of Tor Vergata, Rome

SUMMARY. Introduction. Delusional symptoms are heterogeneous and differentially related to insight, depression and another psychological construct, such as deservedness. In this study we explore models of relationships among these constructs, by hypothesizing that lack of insight could predict depression or paranoia, representing these variables outcome or mediator variables alternatively. **Materials.** We evaluated positive psychotic symptoms, depression, persecution, deservedness and insight in a clinical sample of 81 people with schizophrenia or other psychotic spectrum disorders. Using multiple regression analyses we found 2 significant models. In the first one the lack of insight is negatively associated with depression and negatively related to persecution that, as mediator, is positively associated with depression. In the second model, lack of insight is negatively associated with persecution and negatively related to depression that, as mediator, is positively associated with persecution. **Conclusions.** Persecution and depression could be not only predicted by insight but this prediction could also be mediated by the same variables. In both clinical models found in this study, insight does have a relevant role because the awareness about symptom/illness is crucial to the symptoms change. However, having good insight into schizophrenia can also be paradoxically associated with negative subjective states related to depression (the so-called “insight paradox”).

KEY WORDS: schizophrenia, insight, paranoia, deservedness, PaDS, mediation analysis.

RIASSUNTO. Introduzione. I sintomi deliranti sono eterogenei e differentemente correlati all'insight, alla depressione e a un altro importante costrutto psicologico quale è la “deservedness”. In questo studio esploriamo i modelli relazionali tra questi costrutti, ipotizzando che la mancanza di insight possa prevedere la depressione o la paranoia, rappresentando queste variabili alternativamente come outcome o come mediatore. **Materiali.** Abbiamo valutato i sintomi psicotici positivi, la depressione, la persecuzione, la deservedness e l'insight in un campione clinico di 81 persone con schizofrenia o altri disturbi dello spettro psicotico. Utilizzando un'analisi di regressione multipla abbiamo trovato due modelli significativi. Nel primo la mancanza di insight è negativamente associata alla depressione e negativamente correlata alla persecuzione che, come mediatore, è positivamente associato alla depressione. Nel secondo modello, la mancanza di insight è negativamente associata alla persecuzione e negativamente correlata alla depressione che, come mediatore, è positivamente associata alla persecuzione. **Conclusioni.** La persecuzione e la depressione potrebbero non essere predette dalla sola presenza di insight ma anche dalla reciproca presenza delle due stesse variabili. In entrambi i modelli clinici trovati, l'insight ha un ruolo rilevante perché la consapevolezza circa i sintomi e la malattia è cruciale per il cambiamento dei sintomi stessi. Tuttavia, avere un buon insight nella schizofrenia può anche essere associato paradossalmente a vissuti negativi, legati alla depressione (il cosiddetto “paradosso dell'insight”).

PAROLE CHIAVE: schizofrenia, insight, paranoia, persecuzione, PaDS, mediazione.

INTRODUCTION

More than one century of psychopathological analysis does not sharply delineate clear boundaries between paranoid schizophrenia and other delusional disorders and even within schizophrenia. Delusional symptoms are heterogeneous in nature, development and outcomes as widely investigated from historical, psychometric, psychological psychopathological and philosophical perspectives¹⁻⁵.

A complex relationship with affective and mood symptoms is also reported. In particular, several ambiguities emerge from the distinction between ‘mood congruent’ ver-

sus ‘mood incongruent’ delusions⁶ with low self-esteem and negative expectation^{7,8} associated with delusions.

Since delusions have dimensional⁹ and multidimensional attributions¹⁰, it could be difficult to separate delusions within diagnostic categories but even between mental disorders and reasoning biases in non-clinical sample¹.

A number of models of delusion formation have been elaborated¹¹ including probabilistic reasoning, attributional style, metacognition, self-concepts attribution^{7,12,13} but not clear-cut points of separation between psychiatric diagnoses are feasible^{14,15}.

In the last two decades, it has been theorized how an individual's self-concept, including both self-esteem and self-schemas, low mood and negative schematic beliefs may relate to the formation and maintenance of persecutory delusions¹⁶⁻²¹.

According to some authors by studying self-esteem is even possible to intercept different types of paranoia²² or its dynamic and fluctuating aspects^{23,24}.

A recent systematic review⁷ investigated the degree to which the persecution was thought to be deserved and reported that low self-esteem is associated with higher perceived deservedness of the persecution. Some studies on people with persecutory delusions^{25,26} found negative correlations between perceived deservedness and self-esteem but comparable degrees of non-deserved paranoia patients' self-esteem to those of the healthy sample²⁷.

Interestingly, one study found a similar relation also in non-clinical sample²⁸ and another one found three subtypes of paranoia in healthy delusion-prone people, in line with the severity of depression, self-esteem and social anxiety²⁹.

Chadwik and Trower reported two types of paranoia, 'Bad-Me' (BM) or punishment paranoia and 'Poor-Me' (PM) or persecution paranoia²⁵. According to these authors, people with PM paranoia 'tend to blame others, to see others as bad, and to see themselves as victims'²², as they believe others are plotting to harm them without any justification. People with BM /punishment paranoia, on the other hand, are individuals who 'tend to blame themselves and see themselves as bad, and view others as justifiably punishing them'²². They are more anxious, depressed and have lower self-esteem.

Furthermore, Melo et al.³⁰ evidenced that there is a lack of measures able to intercept the presence of both paranoia and deservedness, administrable to both clinical and not clinical samples; to this purpose they developed the Persecution and Deservedness Scale (PaDS) getting a valid measure of paranoia and deservedness.

We hypothesized that these clinical features could be due to different level of insight that in turn could affect the relationship between depression and persecution³¹⁻³³.

For a better understanding of these relationships we explore models in which lack of insight could predict depression or paranoia with these variables, representing outcome or mediator variables, alternatively.

METHODS

Participants and procedure

The sample of participants in the study included 81 individuals (65.4% males, n. 53; age: 42.40±14.44 years; educational level: 11.49±3.35 years; age at onset: 28.64±13.63 years), who provided complete data on the variables of interest. The sample was drawn from 100 consecutively admitted patients with 19 patients unable to participate for severity or refusal. Inclusion criteria were diagnosis of schizophrenia or other psychotic spectrum disorders, age between 18 and 65, lack of any intellectual disabilities or drugs abuse. All the subjects provided written informed consent after a complete description of the study, were evaluated by two senior psychiatrists and were diagnosed in according with ICD-9 (schizophrenia n. 34; schizoaffective disorder n. 28; unspecified schizophrenia n. 11;

delusional disorder n. 7; schizophreniform disorder n.1). The evaluations were performed before discharge from a hospital admission in a relatively stable clinical condition that allowed the patients to understand and collaborate to the research project.

Measures

Positive and Negative Symptoms Scale (PANSS)³⁴ is a 30-item scale used to evaluate the presence, absence and severity of Positive (P), Negative (N) and General Psychopathology (G) symptoms of schizophrenia. It has seven positive symptom subscale items, seven negative symptom subscale items, and sixteen general psychopathology symptom items. All items are rated on a 7-point scale (1=absent; 7=extreme). Item G12 was used to evaluate lack of insight.

The Persecution and Deservedness Scale (PaDS)³⁰ is a brief measure for assessing both the severity of paranoid thinking and the degree of perceived deservedness of persecution. Two subscales compose it: one measuring persecution and the other the deservedness, where deservedness ratings are orthogonal to persecution ratings. If so, deservedness cannot be scored if the person does not feel to be persecuted.

The persecution subscale (P) has ten statements of persecutory content, some derived from Fenigstein and Venable's scale³⁵ the others from Melo et al.³⁰ experience of clinical practice with paranoid patients, each of which could be scored from 0 to 4 (anchors: 0, 'certainly false'; 1, 'possibly false'; 2, 'unsure'; 3, 'possibly true'; and 4, 'certainly true').

A deservedness item follows each persecution one and should be completed only if the person scores greater than 1 on the related persecution item; as for P, it is scored 0-4. We used the Italian version of the PaDS³⁶.

The Beck Depression Inventory-FS (formerly known as the BDI-PC)³⁷, contains seven cognitive and affective items from the BDI-II to assess depression. Each item rates on a 4-point scale (0-3). It is scored by summing ratings for each item (range 0-21). Items investigate symptoms of sadness, pessimism, past failure, loss of pleasure, self-dislike, self-critical and suicidal thoughts and wishes during the "past 2 weeks including today".

Analysis

Means and standard deviations were reported for descriptive statistics. Pearson r correlation was used where appropriate. Linear regression analyses were conducted to assess each component of the proposed mediation model.

Mediation analyses have been carried out using a SPSS Macro (i.e., PROCESS)³⁸ for observing how an independent variable (in our case the insight, using PANSS G12) affects a second variable (the dependent one such as depression for the first model or the persecution for the second model (Figure 1) via the inclusion of a third variable (the mediator: persecution in the first model – Figure 1a – and depression in the second – Figure 1b). The bootstrapping method with bias-corrected confidence estimates was used^{39,40}. The 95% confidence interval of the indirect effects was obtained with 5000 bootstrap resamples^{41,42}.

RESULTS

Table 1 reports the clinical features of the sample. Table 2 shows the correlations between the clinical variables.

In the first model (Figure 1a), it was found that lack of insight was negatively associated with depression (B=-1.67, t

Paranoia, depression and lack of insight in schizophrenia: a suggestion for a mediation effect

Table 1. Clinical features of the sample (n=81).

	Mean	Standard Deviation
PANSS G12 Lack of insight	3.61	1.52
PANSS Positive score	17.58	6.52
PANSS Negative score	17.74	6.54
PANSS General score(excluding item G12)	34.46	9.22
PANSS Total score	73.40	16.55
BDI-FS Total score	7.28	5.89
PaDS Persecution	23.08	9.08
PaDS Deservedness	7.74	7.60

PANSS: Positive and Negative Symptoms Scale; BDI-FS: Beck Depression Inventory-Fast Screen; PaDS: Persecution and Deservedness Scale

Table 2. Pearson' r correlations among the clinical variables (n=81).

	BDI total score	Persecution total score	Deservedness total score
Persecution tot	.62**		
Deservedness tot	.30*	.38**	
PANSS Pos	-.29*		-.17
PANSS G12	-.43**	-.44**	-.13

PANSS: Positive and Negative Symptoms Scale
 * p<.01; ** p<.0005;
 Only r values >.25 are reported (except for not significant values of deservedness).

(79)=-4.24, p<.0001). It was also found that lack of insight was negatively related to persecution (B=-2.66, t (79)=-4.43, p<.0001). Lastly, results indicated that the mediator, persecution, was positively associated with depression (B=.35, t (79)=5.61, p<.0001). Because both the a-path and b-path were significant, mediation analyses were tested.

Results of the mediation analysis confirmed the mediating role of persecution in the relation between lack of insight and depression (B=-.94; CI=-1.48 to -.51). In addition, results indicated that the direct effect of lack of insight on depression became non-significant (B=-.73, t (79)=-1.97, NS) when controlling for persecution, thus suggesting full mediation. The completely standardized indirect effect size (ab_{cs}) of this model was -.24.

In the second model (Figure 1b), lack of insight was negatively associated with persecution (B=-2.66, t (79)=-4.43, p<.0001). Moreover, lack of insight was negatively related to depression (B=-1.67, t (79)=-4.24, p<.0001). Lastly, results indicated that the mediator, depression, was positively associated with persecution (B=.82, t (79)=5.61, p<.0001). Also in this case, both the a-path and b-path were significant. Results of the mediation analysis confirmed a mediating role of depression in the relation between lack of insight and persecution (B=-1.39; CI=-2.44 to -.67). In addition, results indicated that




Figure 1. Mediation analysis results.

the direct effect of the lack of insight on persecution changed (B=-1.29, t (79)=-2.29, p<.05) when controlling for depression, thus suggesting a high mediation. The completely standardized indirect effect size (ab_{cs}) of this model was -.23.

DISCUSSION

To our knowledge, this is the first study to explore the relationship between paranoia/deservedness construct, lack of insight and depressive symptoms in schizophrenia or other psychotic spectrum disorders exploring these variables as possible mediator in competing models.

As various type and strength of relationships between depression, insight and paranoia have been reported, it is difficult to believe that these relationships would not interact; besides lack of insight partially mediates the relationship between psychotic factors and global functioning⁴³.

We hypothesized a sort of interaction and found that persecution and depression could be not only predicted by insight but this prediction could also be mediated by the same variables.

Differences between the two models reflect clinical complexity. In other words, people reported higher depression because they feel persecuted or feel persecuted because they are depressed? A first comment is that in both cases insight does have a crucial role because the awareness about symptom/illness is crucial to the following symptoms change.

This result could in part support the notion of BM' and PM paranoia theorized by Trower and Chadwick²² and by

Chadwick et al.²⁵. In particular, the one side of the prediction (i.e. BM paranoia) seems to be supported, as we find a correlation between persecution and depression but not between persecution and PANSS positive as expected.

This finding underline the need to further explore the role of insight as a possible ‘psychological cause of deservedness’ while depression could be a ‘psychological consequence’^{32,44}.

Having good insight into schizophrenia can also be paradoxically associated with negative subjective states related to depression, including demoralization, decreased self-esteem, hopelessness and suicidal ideation^{31,45-48}. This so-called ‘insight paradox’ could contribute to the deservedness of persecution.

This finding is in agreement with a recent cluster analytical study where a cluster of patients with most positive insight (i.e. lower PANSS General Psychopathology Item G12 score), shows a more complex pattern, a somewhat ‘paradoxical’ mixture of positive and negative self-reported personal resources (e.g. low self-esteem, low personal strength, higher level of stigma and emotional coping) and clinical features (e.g. high depressive symptoms). Insight into illness may complicate the relationship between self-reported personal and clinical elements of recovery^{45,49-51}. This cluster may identify persons with greater vulnerability to depression but not to psychotic symptoms, probably due to low self-esteem. From this analysis we can establish that ‘deservedness’ could be the other face of insight and depression in schizophrenia. This conceptualization could open new views in cognitive treatment of delusions⁹.

The relationship between paranoia/deservedness, depression and insight should be further explored to identify ‘target’ of tailored psychosocial or pharmacological treatment able to address patient needs and outcome. As depression has been reported to be associated with persecutory delusion, especially if people felt less powerful than their persecutors¹⁷, global insight evaluation could further address psychosocial treatments.

The lack of correlation between PANSS positive score and PaDS deserves further comments. We rely upon the original PANSS scoring reported by Kay et al.³⁴ and the PANSS ‘positive factor’ could unable to catch the complexity of the delusional world⁵². Alternatively, with the term of ‘positive symptoms’ or delusion/paranoia we include different psychopathological constructs⁹, likely heterogeneous clinical phenomena as delusion formation arise from a complex interaction of emotional process, anomalous experience and cognitive bias that could be difficult to separate from not-delusional features². These symptoms could have different relations to insight so that this evaluation is crucial to establish better description and an adequate assessment and treatment within heterogeneity of delusions⁵³.

Conflict of interests: the authors have no conflict of interests to declare.

REFERENCES

1. Bebbington P, Freeman D. Transdiagnostic extension of delusions: schizophrenia and beyond. *Schizophr Bull* 2017; 43: 273-82.
2. Cutting J, Musalek M. The nature of delusion: psychologically explicable? Psychologically inexplicable? Philosophically explicable? Part 2. *Hist Psychiatry* 2016; 27: 21-37.
3. Lewis A. Paranoia and paranoid: a historical perspective. *Psychol Med* 1970; 1: 2-12.
4. Rawlings D, Freeman JL. A questionnaire for the measurement of paranoia/suspiciousness. *Br J Clin Psychol* 1996; 35 (Pt 3): 451-61.
5. Parnanzone S, Serrone D, Rossetti MC, et al. Alterations of cerebral white matter structure in psychosis and their clinical correlations: a systematic review of Diffusion Tensor Imaging studies. *Riv Psichiatr* 2017; 52: 49-66.
6. Kumazaki T. What is a ‘mood-congruent’ delusion? History and conceptual problems. *Hist Psychiatry* 2011; 22: 315-31.
7. Kesting ML, Lincoln TM. The relevance of self-esteem and self-schemas to persecutory delusions: a systematic review. *Compr Psychiatry* 2013; 54: 766-89.
8. Bentall RP, Rowse G, Kinderman P, et al. Paranoid delusions in schizophrenia spectrum disorders and depression: the transdiagnostic role of expectations of negative events and negative self-esteem. *J Nerv Ment Dis* 2008; 196: 375-83.
9. Freeman D, Garety PA. Comments on the content of persecutory delusions: does the definition need clarification? *Br J Clin Psychol* 2000; 39 (Pt 4): 407-14.
10. Freeman D. Suspicious minds: the psychology of persecutory delusions. *Clin Psychol Rev* 2007; 27: 425-57.
11. Bell V, Halligan PW, Ellis HD. Explaining delusions: a cognitive perspective. *Trends Cogn Sci* 2006; 10: 219-26.
12. Morrison AP, Shryane N, Fowler D, et al. Negative cognition, affect, metacognition and dimensions of paranoia in people at ultra-high risk of psychosis: a multi-level modelling analysis. *Psychol Med* 2015; 45: 2675-84.
13. Tiernan B, Tracey R, Shannon C. Paranoia and self-concepts in psychosis: a systematic review of the literature. *Psychiatry Res* 2014; 216: 303-13.
14. Monti MR, Stanghellini G. Psychopathology: an edgeless razor? *Compr Psychiatry* 1996; 37: 196-204.
15. Stanghellini G, Rossi R. Pheno-phenotypes: a holistic approach to the psychopathology of schizophrenia. *Curr Opin Psychiatry* 2014; 27: 236-41.
16. Freeman D, Garety PA, Bebbington PE, et al. Psychological investigation of the structure of paranoia in a non-clinical population. *Br J Psychiatry* 2005; 186: 427-35.
17. Green C, Garety PA, Freeman D, et al. Content and affect in persecutory delusions. *Br J Clin Psychol* 2006; 45: 561-77.
18. Krabbendam L, Janssen I, Bak M, et al. Neuroticism and low self-esteem as risk factors for psychosis. *Soc Psychiatry Psychiatr Epidemiol* 2002; 37: 1-6.
19. Martin JA, Penn DL. Social cognition and subclinical paranoid ideation. *Br J Clin Psychol* 2001; 40: 261-5.
20. Bucci P, Galderisi S, Mucci A, et al. Premorbid academic and social functioning in patients with schizophrenia and its associations with negative symptoms and cognition. *Acta Psychiatr Scand* 2018; 138: 253-66.
21. Rocca P, Galderisi S, Rossi A, et al. Disorganization and real-world functioning in schizophrenia: Results from the multicenter study of the Italian Network for Research on Psychoses. *Schizophr Res* 2018; 201: 105-12.
22. Trower P, Chadwick P. Pathways to defense of the Self: a theory of two types of paranoia. 1995; 2: 263-78.
23. Bentall RP, Corcoran R, Howard R, Blackwood N, Kinderman P. Persecutory delusions: a review and theoretical integration. *Clin Psychol Rev* 2001; 21: 1143-92.
24. Bentall RP, Kinderman P, Kaney S. The self, attributional processes and abnormal beliefs: towards a model of persecutory delusions. *Behav Res Ther* 1994; 32: 331-41.
25. Chadwick PD, Trower P, Juusti-Butler TM, Maguire N. Phenomenological evidence for two types of paranoia. *Psychopathology* 2005; 38: 327-33.

Paranoia, depression and lack of insight in schizophrenia: a suggestion for a mediation effect

26. Fowler D, Hodgekins J, Garety P, et al. Negative cognition, depressed mood, and paranoia: a longitudinal pathway analysis using structural equation modeling. *Schizophr Bull* 2012; 38: 1063-73.
27. Fornells-Ambrojo M, Garety PA. Understanding attributional biases, emotions and self-esteem in 'poor me' paranoia: findings from an early psychosis sample. *Br J Clin Psychol* 2009; 48: 141-62.
28. Pickering L, Simpson J, Bentall RP. Insecure attachment predicts proneness to paranoia but not hallucinations. *Personality and Individual Differences* 2008; 44: 1212-24.
29. Combs DR, Penn DL, Chadwick P, et al. Subtypes of paranoia in a nonclinical sample. *Cogn Neuropsychiatry* 2007; 12: 537-53.
30. Melo S, Corcoran R, Shryane N, Bentall RP. The persecution and deservedness scale. *Psychol Psychother* 2009; 82: 247-60.
31. Lysaker PH, Vohs J, Hasson-Ohayon I, et al. Depression and insight in schizophrenia: comparisons of levels of deficits in social cognition and metacognition and internalized stigma across three profiles. *Schizophr Res* 2013; 148: 18-23.
32. Rossi A, Galderisi S, Rocca P, et al. Personal resources and depression in schizophrenia: the role of self-esteem, resilience and internalized stigma. *Psychiatry Res* 2017; 256: 359-64.
33. Belvederi Murri M, Respino M, Innamorati M, et al. Is good insight associated with depression among patients with schizophrenia? Systematic review and meta-analysis. *Schizophr Res* 2015; 162: 234-47.
34. Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull* 1987; 13: 261-76.
35. Fenigstein A, Venable PA. Paranoia and self-consciousness. *J Pers Soc Psychol* 1992; 62: 129-38.
36. Serrone D, Stratta P, Riccardi I, et al. Is persecution deserved? A study on a non-clinical Italian sample using Persecution and Deservedness Scale (PaDS). *J Psychopathol* 2018; 24: 24-30.
37. Beck AT, Steer RA, Brown GK. BDI - FastScreen for Medical Patients. San Antonio (TX): The Psychological Corporation, 2000.
38. Hayes AF. Introduction to mediation, moderation and conditional process analysis. New York: The Guilford Press, 2013.
39. Mackinnon DP, Lockwood CM, Williams J. Confidence limits for the indirect effect: distribution of the product and resampling methods. *Multivariate Behav Res* 2004; 39: 99.
40. Preacher KJ, Hayes AF. SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behav Res Methods Instrum Comput* 2004; 36: 717-31.
41. Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* 2008; 40: 879-91.
42. Preacher KJ, Kelley K. Effect size measures for mediation models: quantitative strategies for communicating indirect effects. *Psychol Methods* 2011; 16: 93-115.
43. Quartini A, Pacitti F, Bersani G, Iannitelli A. From adolescent neurogenesis to schizophrenia: Opportunities, challenges and promising interventions. *Biomedical Reviews* 2017; 28: 66-73.
44. Rossi A, Amore M, Galderisi S, et al. The complex relationship between self-reported 'personal recovery' and clinical recovery in schizophrenia. *Schizophr Res* 2018; 192: 108-12.
45. Belvederi Murri M, Amore M, Calcagno P, et al. The "Insight Paradox" in schizophrenia: magnitude, moderators and mediators of the association between insight and depression. *Schizophr Bull* 2016; 42: 1225-33.
46. Lysaker PH, Pattison ML, Leonhardt BL, Phelps S, Vohs JL. Insight in schizophrenia spectrum disorders: relationship with behavior, mood and perceived quality of life, underlying causes and emerging treatments. *World Psychiatry* 2018; 17: 12-23.
47. Sampogna G, Fiorillo A, Luciano M, et al. A randomized controlled trial on the efficacy of a psychosocial behavioral intervention to improve the lifestyle of patients with severe mental disorders: study protocol. *Front Psychiatry* 2018; 9: 235.
48. Biondi M, Iannitelli A, Ferracuti S. [On the unpredictability of suicide]. *Riv Psichiatr* 2016; 51: 167-71.
49. Lysaker PH, Roe D, Yanos PT. Toward understanding the insight paradox: internalized stigma moderates the association between insight and social functioning, hope, and self-esteem among people with schizophrenia spectrum disorders. *Schizophr Bull* 2007; 33: 192-9.
50. Lysaker PH, Vohs J, Hillis JD, et al. Poor insight into schizophrenia: contributing factors, consequences and emerging treatment approaches. *Expert Rev Neurother* 2013; 13: 785-93.
51. Iannitelli A, Parnanzone S, Pizziconi G, Riccobono G, Pacitti F. Psychodynamically oriented psychopharmacotherapy: towards a necessary synthesis. *Front Hum Neurosci* 2019; 13: 15.
52. Mishara AL, Fusar-Poli P. The phenomenology and neurobiology of delusion formation during psychosis onset: Jaspers, Truman symptoms, and aberrant salience. *Schizophr Bull* 2013; 39: 278-86.
53. Kimhy D, Goetz R, Yale S, Corcoran C, Malaspina D. Delusions in individuals with schizophrenia: factor structure, clinical correlates, and putative neurobiology. *Psychopathology* 2005; 38: 338-44.